arXiv:1209.0660v1 [math.RA] 4 Sep 2012

Commuting normal idempotent tropical matrices: an algebraic–geometric approach

J. Linde M.J. de la Puente[*]

AMS class.: 15A80; 52B11; 14T05.

Keywords and phrases: commuting matrices, normal matrix, idempotent matrix, tropical simplex, convexity.

Abstract

We address two related questions. Let two real square normal tropically idempotent matrices A, B be given. Question 1: find conditions for A, B to commute under tropical multiplication. Question 2: prove that the intersection of the two tropical simplexes $\mathcal{T}_A, \mathcal{T}_B$ is a tropical simplex.

1 Introduction and notations

Write $\oplus = \max$ and $\odot = +$. These are the tropical operations addition and multiplication. Write $\overline{\mathbb{R}} := \mathbb{R} \cup \{-\infty\}$. For $n \in \mathbb{N}$, set $[n] := \{1, 2, \ldots, n\}$. Let $\mathbb{R}^{n\times n}$ denote the set of $n \times n$ real matrices, and $\overline{\mathbb{R}}^{n\times n}$ the set of $n \times n$ matrices with entries in $\overline{\mathbb{R}}$ *and, at least, one real entry in each row and column*. Define tropical sum and product of matrices following the same rules of classical linear algebra, but replacing addition (multiplication) by tropical addition (multiplication). We will never use classical sum or multiplication of matrices, in this note. $A \odot B$ will be written AB, for simplicity, for matrices A, B.

Normal idempotent matrices (NI, for short) are characterized by n equalities and $n(n-1)^2$ linear inequalities. The class of NI matrices is interesting because any square real matrix A can be assumed to be normal, up to coordinate changes. Moreover, for any real normal order n matrix A, the tropical power A^{n-1} is NI, and A^{n-1} carries most of the information of A. NI matrices are increasing, strongly definite, visualized, and Kleene stars (see [5, 6, 8, 30]). We do not use the digraph associated to a matrix or eigenvectors and eigenvalues, when dealing with NI matrices. Our basic technique is handling inequalities.

The geometric counterpart of a NI matrix in the tropical projective space is a tropical simplex which contains the origin and, *moreover*, is an alcoved polytope

---

[*] Partially supported by UCM research group 910444. Corresponding author

i.e., a convex set defined by real linear inequalities of the sort $p_i \leq x_i \leq q_i$ and $p_{ij} \leq x_i - x_j \leq q_{ij}$, some $p_i, q_i, p_{ij}, q_{ij} \in \mathbb{R}$ (see [18, 19, 21, 22, 31]).

The problem with NI matrices is that this class is not closed under tropical sum or (left or right) multiplication. However, *if A, B are NI and AB, BA are also NI, then A and B commute and the intersection $\mathcal{T}_A \cap \mathcal{T}_B$ is a tropical simplex. If the tropical sum $A \oplus B$ is NI, the same holds.* This is essentially what we prove in this note. Our main results are theorems 3, 4, 9 and 10. A key role is played by lemma 1 and theorem 2.

Basic facts about tropical algebra and geometry and variations of them (also called max–plus, max–linear, idempotent mathematics, etc.) can be found in the following texts: [1, 3, 5, 7, 8, 10, 12, 13, 14, 23, 24, 26, 29, 33]. Convexity of sets tropically spanned by finitely many elements has been studied in [2, 6, 9, 11, 15, 16, 17, 18, 19, 30].

The $n \times n$ matrix I, having null diagonal and $-\infty$ elsewhere, is an identity for tropical multiplication. 0 will denote the $n \times n$ null matrix. For $d_1, \ldots, d_n \in \mathbb{R}$, $\operatorname{diag}(d_1, \ldots, d_n)$ is the $n \times n$ matrix having $d_1, \ldots, d_n$ on the diagonal and $-\infty$ elsewhere.

2 Normal and normal idempotent matrices

By definition, a square matrix A is *normal* if $a_{ii} = 0$ and $a_{ij} \leq 0$, all i, j. Equivalently, $I \leq A \leq 0$. Notice that $A0 = 0 = 0A$, when A is normal. Since tropical (left or right) multiplication is monotonic, then $I \leq A \leq A^2 \leq \ldots \leq A^{n-1} \leq A^n \leq A^{n+1} \leq \ldots \leq 0$, if A is normal. A theorem of Yoeli's (see [32]) says that this sequence stabilizes: $A^{n-1} = A^n$.

Given $A = (a_{ij})$, $B = (b_{ij})$ normal matrices, set

- $M := A \oplus B$, the *maximum matrix*, $M = (m_{ij})$,
- $N := AB \oplus BA$, the *maximum of the products*,
- $H := \min\{AB, BA\}$, the *minimum of the products*,
- $c_{ij}(k) := a_{ik} + b_{kj}$, $C(k) := (c_{ij}(k))$, with $i, j, k \in [n]$, so that
$$AB = \bigoplus_{k \in [n]} C(k) \oplus A \oplus B,$$
- $d_{ij}(k) := b_{ik} + a_{kj}$, $D(k) := (d_{ij}(k))$, with $i, j, k \in [n]$, so that
$$BA = \bigoplus_{k \in [n]} D(k) \oplus A \oplus B.$$

By the *Hungarian method*, (see [5, 20, 27]), for any matrix $A \in \mathbb{R}^{n\times n}$, there exist (not necessarily unique) matrices $P, Q \in \overline{\mathbb{R}}^{n\times n}$ and $N \in \mathbb{R}^{n\times n}$ such that $N = PAQ$, with N normal. The matrices P and Q are generalized permutation matrices. This means that P is obtained from a diagonal matrix by simultaneous permutation of rows and columns (same for Q). On the tropical projective space (see definition on p. 6), the matrix P acts as a change of coordinates (a translation and a relabeling of coordinates), and the matrix Q acts on A by relabeling its columns. Therefore, A can be assumed to be normal, up to coordinate changes.

A matrix A is *idempotent* if $A^2 = A$. A *normal idempotent* matrix (NI, for short) A satisfies

$$a_{ii} = 0, \quad a_{ij} \le 0 \text{ and } a_{ik} + a_{kj} \le a_{ij}, \quad i, j, k \in [n]. \tag{1}$$

These are n equalities and $n(n-1)+n(n-1)(n-2) = n(n-1)^2$ linear inequalities. The *tropical determinant* of A is defined as

$$|A|_{\text{trop}} = \max_{\sigma\in S_n}\{a_{1\sigma(1)} + a_{2\sigma(2)} + \cdots + a_{n\sigma(n)}\},$$

where S_n denotes the permutation group in n symbols. It is also called tropical permanent. A is *tropically regular* (resp. *tropically singular*) if this maximum is attained only once (resp. at lest twice). Strongly regular is another denomination for tropical regular. In [25] it is proved that if AB is tropically regular, then A and B are tropically regular and

$$|AB|_{\text{trop}}| = |A|_{\text{trop}} + |B|_{\text{trop}}. \tag{2}$$

From now on, $A, B \in \mathbb{R}^{n\times n}$.

Here is a list of immediate properties, some true for normal matrices, and some true for NI matrices.

1. If A, B are normal, then AB, BA and M, N, H are normal.
2. If A, B are normal, then $M \le H$.

   Indeed, $AB = \bigoplus_k C(k) \oplus M$, $BA = \bigoplus_k D(k) \oplus M$, whence $M \le AB$ and $M \le BA$. An alternative proof is the following: $B = IB \le AB$, $B = BI \le BA$, $A = IA \le BA$, $A = AI \le AB$, by monotonicity of multiplication, whence $M \le H$.
3. If A, B are NI, then AB, BA or M may fail to be NI.

   Example 1: A, B are NI but $M \ne AB \ne BA \ne M$ are not, with

$$A = \begin{bmatrix} 0 & -3 & -3 \\ 0 & 0 & 0 \\ 0 & -3 & 0 \end{bmatrix}, \quad B = \begin{bmatrix} 0 & -2 & -3 \\ -2 & 0 & -1 \\ -1 & -3 & 0 \end{bmatrix}$$

$$AB = \begin{bmatrix} 0 & -2 & -3 \\ 0 & 0 & 0 \\ 0 & -2 & 0 \end{bmatrix}, BA = \begin{bmatrix} 0 & -2 & -2 \\ 0 & 0 & 0 \\ 0 & -3 & 0 \end{bmatrix}, M = \begin{bmatrix} 0 & -2 & -3 \\ 0 & 0 & 0 \\ 0 & -3 & 0 \end{bmatrix}.$$

Example 2: A, B and $AB = BA$ are NI but M is not, with

$$A = \begin{bmatrix} 0 & -4 & -6 & -3 \\ -6 & 0 & -4 & -3 \\ -3 & -6 & 0 & -3 \\ -6 & -3 & -3 & 0 \end{bmatrix}, \quad B = \begin{bmatrix} 0 & -4 & -4 & -6 \\ -2 & 0 & -3 & -4 \\ -5 & -6 & 0 & -5 \\ -6 & -5 & -2 & 0 \end{bmatrix}$$

$$AB = BA = M^2 = \begin{bmatrix} 0 & -4 & -4 & -3 \\ -2 & 0 & -3 & -3 \\ -3 & -6 & 0 & -3 \\ -5 & -3 & -2 & 0 \end{bmatrix}, \quad M = \begin{bmatrix} 0 & -4 & -4 & -3 \\ -2 & 0 & -3 & -3 \\ -3 & -6 & 0 & -3 \\ -6 & -3 & -2 & 0 \end{bmatrix}.$$

4. If A, B and M are NI, then $AB = BA$ (and, by theorem 3 below, AB is NI).

   Indeed, $M = M^2 = (A \oplus B)^2 = A^2 \oplus B^2 \oplus N = A \oplus B \oplus N = M \oplus N$, whence $N \leq M$ and $M \leq H \leq N$ by item 2, showing $M = H = N$; in particular $AB = BA$.

Lemma 1. *If A, B are normal, then*

1. $ABA = A^2 \oplus N$,
2. $BAB = B^2 \oplus N$,
3. $A^2B = A^2 \oplus AB$,
4. $B^2A = B^2 \oplus BA$,
5. $BA^2 = A^2 \oplus BA$,
6. $AB^2 = B^2 \oplus AB$.

Proof. To prove (1), fix $i \neq j \in [n]$.

$$(ABA)_{ij} = \max_{k \in [n]} \{(AB)_{ik} + a_{kj}\} = \max_{k \in [n]} \{\max_{l \in [n]} \{a_{il} + b_{lk}\} + a_{kj}\} = \max_{k,l \in [n]} \{a_{il} + b_{lk} + a_{kj}\}.$$

If two indices coincide, the corresponding summand vanishes; otherwise, the summand is non–positive. Therefore, summands having three non–positive terms are irrelevant and

$$(ABA)_{ij} = \max_{k \in [n]} \{a_{ij}, b_{ij}, a_{ik} + a_{kj}, c_{ij}(k), d_{ij}(k)\}.$$

Using $AB = \bigoplus_k C(k) \oplus A \oplus B$ and $BA = \bigoplus_k D(k) \oplus A \oplus B$, we get $ABA = A^2 \oplus AB \oplus BA$.

To prove (3), fix $i \neq j \in [n]$.

$$(A^2B)_{ij} = \max_{k\in[n]}\{(A^2)_{ik}+b_{kj}\} = \max_{k\in[n]}\{\max_{l\in[n]}\{a_{il}+a_{lk}\}+b_{kj}\} = \max_{k,l\in[n]}\{a_{il}+a_{lk}+b_{kj}\}.$$

Thus

$$(A^2B)_{ij} = \max_{k\in[n]}\{a_{ij}, b_{ij}, a_{ik}+a_{kj}, c_{ij}(k)\},$$

whence $A^2B = A^2 \oplus AB$.

The other items are proved similarly. □

Theorem 2. *If A, B are normal, then $(AB)^2 = (BA)^2$.*

Proof. $(AB)^2 = (ABA)B = (A^2 \oplus AB \oplus BA)B = A^2B \oplus AB^2 \oplus BAB = A^2 \oplus B^2 \oplus N$, using lemma 1. Similarly, we get $(BA)^2 = A^2 \oplus B^2 \oplus N$. □

3 Commuting NI matrices

Theorem 3. *If A, B are NI, then A and B commute if and only if AB and BA are idempotent.*

Proof. If A and B commute, then $(AB)^2 = A^2B^2 = AB$ and $(BA)^2 = B^2A^2 = BA$. Conversely, if AB and BA are idempotent, then $AB = (AB)^2 = (BA)^2 = BA$, by the previous theorem. □

Theorem 4. *If A, B are NI of order 3, then A and B commute if and only if $AB = BA = M$.*

Proof. Let us derive a contradiction from $AB = BA \neq M$. Write $C = (c_{ij})$, $D = (d_{ij})$ with $c_{ij} = a_{ik} + b_{kj}$, $d_{ij} = b_{ik} + a_{kj}$, $\{i, j, k\} = [3]$. We have $AB = M \oplus C = M \oplus D = BA$. There exist $i \neq j \in [3]$ such that $m_{ij} < c_{ij}$. Say $i = 1$, $j = 2$. Then $a_{12} < a_{13} + b_{32}$ and $b_{12} < a_{13} + b_{32}$. Since A, B are NI, then $a_{32} \leq a_{12} - a_{13} < b_{32}$ and $b_{13} \leq b_{12} - b_{32} < a_{13}$. Adding up, we get $b_{13} + a_{32} < a_{13} + b_{32}$, meaning that $d_{12} < c_{12}$, whence $(AB)_{12} \neq (BA)_{12}$. □

It is good now to go back to examples 1 and 2.

Here is an easy way to produce NI matrices which commute.

Lemma 5. *For each negative real number m, any two matrices A, B having null diagonal and all off–diagonal entries in the closed interval $[2m, m]$ are NI and satisfy $AB = BA = M$.*

Proof. Let $a_{ii} = b_{ii} = 0$ and $2m \leq a_{ij}, b_{ij} \leq m$. Then $a_{ik} + a_{kj} \leq 2m \leq a_{ij}$, whence A is NI. Same for B.

Now, fix $i \neq j \in [n]$. We have $c_{ij}(k) \leq 2m \leq a_{ij}$, whence

$$(AB)_{ij} = \max_{k\in[n]}\{a_{ij}, b_{ij}, c_{ij}(k)\} = \max_{k\in[n]}\{a_{ij}, b_{ij}\} = (A \oplus B)_{ij}.$$

Thus, $AB = M$. Similarly, we get $BA = M$. ☐

4 The algebraic–geometric interplay

The projective tropical $n-1$–dimensional space, denoted $\mathbb{TP}^{n-1}$ is, in this note, the quotient space $\overline{\mathbb{R}}^n \setminus \{(-\infty, \dots, -\infty)\}/\sim$, where $(a_1, \dots, a_n) \sim (b_1, \dots, b_n)$ if $(a_1, \dots, a_n) = (\lambda + b_1, \dots, \lambda + b_n)$, for some $\lambda \in \mathbb{R}$. The equivalence class of $(a_1, \dots, a_n)$ will be denoted $[a_1, \dots, a_n]$. Let coordinates on $\mathbb{TP}^{n-1}$ be denoted by $X_1, \dots, X_n$. For any real point in $\mathbb{TP}^{n-1}$, we can choose a representative having null last coordinate $(a_1, \dots, a_{n-1}, 0)$. In this case, we say that we *work in* $X_n = 0$. This allows us to identify the real part of $\mathbb{TP}^{n-1}$ with $\mathbb{R}^{n-1}$.

From now on, we will write points in $\mathbb{TP}^{n-1}$ as columns.

By definition, a *tropical simplex* in $\mathbb{TP}^{n-1}$ is the set of points tropically spanned by n given real points. We arrange the coordinates of the points as the columns of an $n \times n$ real matrix A and denote the simplex by $\mathcal{T}_A$ (in the literature other names are used, such as tropically convex set, column space of A, etc.). In other words, $\mathcal{T}_A$ is the image of the map $f_A : \mathbb{TP}^{n-1} \to \mathbb{TP}^{n-1}$ given by $x \mapsto Ax$. In the tropical projective space, $\mathcal{T}_A$ is compact, connected but not convex, in general (compactness does not hold in the tropical affine space!). The topological dimension of $\mathcal{T}_A$ is $\leq n-1$ with equality if and only if A is tropically regular. Equivalently, $\dim \mathcal{T}_A = n-1$ if and only if the columns of A are not all contained in one tropical hyperplane, meaning that there is no $p = [p_1, \dots, p_n]^t \in \mathbb{TP}^{n-1}$ such that for each $j \in [n]$, the maximum

$$\max\{p_1 + a_{1j}, \dots, p_n + a_{nj}\} \tag{3}$$

is attained at least twice; see [4, 5, 15, 16, 17, 29]. Generically, $\mathcal{T}_A$ is the union of one cell of dimension $n-1$ and finitely many cells of dimensions m, for $0 \leq m \leq n-2$; see [9].

To sketch the simplex $\mathcal{T}_A$ in $\mathbb{R}^{n-1}$, we use the matrix $A_0 = (\alpha_{ij})$, with $\alpha_{ij} = a_{ij} - a_{nj}$. For instance, (see figure 1) in example 1 we have

$$A_0 = \begin{bmatrix} 0 & 0 & -3 \\ 0 & 3 & 0 \\ 0 & 0 & 0 \end{bmatrix}, \quad B_0 = \begin{bmatrix} 1 & 1 & -3 \\ -1 & 3 & -1 \\ 0 & 0 & 0 \end{bmatrix}$$

$$(AB)_0 = \begin{bmatrix} 0 & 0 & -3 \\ 0 & 2 & 0 \\ 0 & 0 & 0 \end{bmatrix}, (BA)_0 = \begin{bmatrix} 0 & 1 & -2 \\ 0 & 3 & 0 \\ 0 & 0 & 0 \end{bmatrix}, M_0 = \begin{bmatrix} 0 & 1 & -3 \\ 0 & 3 & 0 \\ 0 & 0 & 0 \end{bmatrix}.$$



Figure 1: Tropical triangles corresponding to example 1. The origin in $\mathbb{R}^2$ is in red, $\mathcal{T}_A$ in magenta, $\mathcal{T}_B$ in blue.

In example 2, we get

$$A_0 = \begin{bmatrix} 6 & -1 & -3 & -3 \\ 0 & 3 & -1 & -3 \\ 3 & -3 & 3 & -3 \\ 0 & 0 & 0 & 0 \end{bmatrix}, \quad B_0 = \begin{bmatrix} 6 & 1 & -2 & -6 \\ 4 & 5 & -1 & -4 \\ 1 & -1 & 2 & -5 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

$$(AB)_0 = \begin{bmatrix} 5 & -1 & -2 & -3 \\ 3 & 3 & -1 & -3 \\ 2 & -3 & 2 & -3 \\ 0 & 0 & 0 & 0 \end{bmatrix}, \quad M_0 = \begin{bmatrix} 6 & -1 & -2 & -3 \\ 4 & 3 & -1 & -3 \\ 3 & -3 & 2 & -3 \\ 0 & 0 & 0 & 0 \end{bmatrix}.$$

Some computations provide the following information. The tropical tetrahedra $\mathcal{T}_A, \mathcal{T}_B, \mathcal{T}_{AB}$ are convex and

- The f–vector of $\mathcal{T}_A$ is $(14, 21, 9)$. It has 4 quadrilateral, 4 pentagonal and one hexagonal facets.
- The f–vector of $\mathcal{T}_B$ is $(20, 30, 12)$. It has 4 quadrilateral, 4 pentagonal and 4 hexagonal facets.
- The f–vector of $\mathcal{T}_{AB}$ is $(15, 23, 10)$. It has one triangular, 3 quadrilateral, 5 pentagonal and one hexagonal facets. See figure 2.



Figure 2: Tropical tetrahedra corresponding to example 2: $\mathcal{T}_A$ in magenta, $\mathcal{T}_B$ in blue, $\mathcal{T}_{AB}$ in black.

However, $\mathcal{T}_M$ is not convex since $[6,4,3,0]^t$ and $[-1,3,-3,0]^t \in \mathcal{T}_M$ but their mid–point is not in $\mathcal{T}_M$.

Consider the following open cells in $\mathbb{R}^{n-1}$: $S_n = \{0 < x_k : k \in [n-1]\}$ and $S_j = \{0 < x_j \text{ and } x_k < x_j : k \in [n-1]\}$, for $j \in [n-1]$. The geometric meaning of normality of A is:

$$\operatorname{col}(A_0, j) \in \overline{S_j}, \quad \text{for all} \quad j \in [n] \tag{4}$$

where the bar denotes usual closure. Notice that $\bigcup_{j\in[n]} \overline{S_j}$ is the whole $\mathbb{R}^{n-1}$.



Figure 3: Tropical triangles corresponding to example 3. The origin in $\mathbb{R}^2$ is in red, $\mathcal{T}_A$ in magenta, $\mathcal{T}_B$ in blue.

Here is a list of properties and remarks:

1. A, B normal and $A \leq B \not\Rightarrow \mathcal{T}_A \supseteq \mathcal{T}_B$, in general.

   Example 3:

$$A = \begin{bmatrix} 0 & 0 & -2 \\ -1 & 0 & -3 \\ -2 & 0 & 0 \end{bmatrix}, \quad B = \begin{bmatrix} 0 & 0 & -1 \\ -1 & 0 & -3 \\ -2 & 0 & 0 \end{bmatrix}$$

$$A_0 = \begin{bmatrix} 2 & 0 & -2 \\ 1 & 0 & -3 \\ 0 & 0 & 0 \end{bmatrix}, \quad B_0 = \begin{bmatrix} 2 & 0 & -1 \\ 1 & 0 & -3 \\ 0 & 0 & 0 \end{bmatrix}$$

   and $[-1, -3, 0]^t \in \mathcal{T}_B \setminus \mathcal{T}_A$; see figure 3.

2. If A is normal, then $\mathcal{T}_A \supseteq \mathcal{T}_{A^2} \supseteq \ldots \supseteq \mathcal{T}_{A^{n-1}} = \mathcal{T}_{A^n}$; see [4, 5, 25, 28].

3. If A is normal, then $\mathcal{T}_{A^{n-1}}$ is the set of fixed points of f_A; see [4, 5, 25, 28]. Indeed, for some $y \in \mathbb{TP}^{n-1}$, we have $x = A^{n-1}y = A^n y = Ax$.

4. If A is normal, then the origin lies in $\mathcal{T}_{A^{n-1}}$.

Any zero–diagonal matrix A in $\mathbb{R}^{n\times n}$ induces the following convex (possibly empty!) polytope in $\mathbb{R}^{n-1}$

$$\mathcal{C}_A := \left\{ x \in \mathbb{R}^{n-1} : \begin{array}{c} a_{in} \leq x_i \leq -a_{ni} \\ a_{ik} \leq x_i - x_k \leq -a_{ki} \end{array} ;\ i, k \in [n-1] \right\}. \tag{5}$$

Theorem 6. *[18] If A is normal, then $\mathcal{C}_A \subseteq \mathcal{T}_A$. Moreover, A is NI if and only if $\mathcal{C}_A = \mathcal{T}_A$.*

We give the proof below, for completeness.

Proof. Recall the definition of α_{ij} in p.6. For $i, j, k, l \in [n]$, we have

1. $\alpha_{ni} = 0$ and $\alpha_{ii} = -a_{ni}$,

2. $\alpha_{in} = a_{in}$,

3. $\alpha_{ij} - \alpha_{jj} = a_{ij}$.

Assuming A is NI, then

4. $a_{in} \leq \alpha_{ij} \leq -a_{ni}$,

5. $a_{ik} \leq \alpha_{ij} - \alpha_{kj} = a_{ij} - a_{kj} \leq -a_{ki}$.

Given a point $x \in \mathcal{C}_A$, consider scalars $\lambda_i = x_i + a_{ni} \leq 0$, for $i \in [n-1]$ and write $\lambda = [\lambda_1, \ldots, \lambda_{n-1}, 0]^t$. Then $[x_1, \ldots, x_{n-1}, 0]^t = A_0 \lambda$ (this is a statement about n maxima, which holds due to the $2(n-1)^2$ inequalities defining $\mathcal{C}_A$), proving that $x \in \mathcal{T}_A$. This proves that $\mathcal{C}_A \subseteq \mathcal{T}_A$.

Next we prove that, for every x, y in $\mathcal{C}_A$, the tropical segment joining x and y, denoted $\mathcal{T}_{x,y}$, is contained in $\mathcal{C}_A$. Call this *fact 1*. Indeed: a point z in $\mathcal{T}_{x,y}$ has coordinates $z_n = 0 = \max\{\lambda, \mu\}$ and

$$z_i = \max\{\lambda + x_i, \mu + y_i\},$$

for $i \in [n-1]$ and some $\lambda, \mu \in \mathbb{R}$. Say $\lambda = 0, \mu \leq 0$; then

$$a_{in} \leq x_i \leq \max\{x_i, \mu + y_i\} = z_i \leq \max\{x_i, y_i\} \leq -a_{ni},$$

for $i \in [n-1]$. Choosing now $i, k \in [n-1]$, we want to prove that $a_{ik} \leq z_i - z_k \leq -a_{ki}$.

If $x_i = z_i$ and $x_k = z_k$ or $\mu + y_i = z_i$ and $\mu + y_k = z_k$, $a_{ik} \leq z_i - z_k \leq -a_{ki}$ is obvious.

If $\mu + y_i = z_i$ and $x_k = z_k$, then $a_{ik} \leq x_i - x_k \leq z_i - z_k = \mu + y_i - x_k \leq \mu + y_i - \mu - y_k = y_i - y_k \leq -a_{ki}$. If $\mu + y_k = z_k$ and $x_i = z_i$, then the situation is similar. This shows that $\mathcal{T}_{x,y} \subseteq \mathcal{C}_A$.

Now we are ready to prove $\mathcal{C}_A \supseteq \mathcal{T}_A$, if A is NI.

First, notice that each column of A_0 belongs to $\mathcal{C}_A$, by items 4 and 5 above. Call this *fact 2*.

Any point y in $\mathcal{T}_A$ admits coordinates of the form $A_0\lambda$, for some $\lambda = [\lambda_1, \ldots, \lambda_n]^t$, with $y_n = \max\{\lambda_1, \ldots, \lambda_n\} = 0$. Then $\lambda_j = 0$, for some j. Say $j = 1$ and write $y = x \oplus \lambda_3 + \operatorname{col}(A_0, 3) \oplus \cdots \oplus \lambda_n + \operatorname{col}(A_0, n)$, with $x = \operatorname{col}(A_0, 1) \oplus \lambda_2 + \operatorname{col}(A_0, 2)$. Now x belongs to $\mathcal{C}_A$, by facts 1 and 2. Then, by fact 1, in finitely many steps, we show that $y \in \mathcal{C}_A$, proving that $\mathcal{T}_A \subseteq \mathcal{C}_A$. ☐

Here is a list of properties and remarks:

1. If A is NI, then A and $\mathcal{T}_A$ determine each other.

   Indeed, we can recover A from $\mathcal{T}_A$, because the NI hypothesis implies that $\mathcal{T}_A = \mathcal{C}_A$ is an alcoved polytope in $\mathbb{R}^{n-1}$ containing the origin. For $i \neq j$, a_{ij} is directly read from $\mathcal{C}_A$. Then set $A = (a_{ij})$, with $a_{ii} = 0$.

   Example 4: given $\mathcal{C} = \{0 \leq x \leq 4,\ 0 \leq y \leq 2,\ -1 \leq x - y\}$, (see figure 4) we notice that the maximum of $x - y$ over $\mathcal{C}$ is 4, whence $\mathcal{C} = \mathcal{C}_A$ with

$$A = \begin{bmatrix} 0 & -1 & 0 \\ -4 & 0 & 0 \\ -4 & -2 & 0 \end{bmatrix}, \quad A_0 = \begin{bmatrix} 4 & 1 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 0 \end{bmatrix}.$$

Alternatively, from $\mathcal{C}$ we write the auxiliary matrix

$$A' = \begin{bmatrix} 0 & -1 & 0 \\ -\infty & 0 & 0 \\ -4 & -2 & 0 \end{bmatrix}, \quad A = (A')^{n-1} = (A')^2.$$



Figure 4: Tropical triangle corresponding to example 4.

2. If A, B are normal, then $A \leq B \Leftrightarrow \mathcal{C}_A \supseteq \mathcal{C}_B$. In particular, $\mathcal{C}_A \cap \mathcal{C}_B = \mathcal{C}_M$.

   This is immediate.

3. If A, B are NI, then $A \leq B \Leftrightarrow \mathcal{T}_A \supseteq \mathcal{T}_B$.

   This is a consequence of theorem 6 and item 2.

4. If A, B are NI, then $\mathcal{T}_A \cap \mathcal{T}_B \subseteq \mathcal{T}_M$.

   Indeed, $\mathcal{T}_A = \mathcal{C}_A$, $\mathcal{T}_B = \mathcal{C}_B$ and $\mathcal{C}_A \cap \mathcal{C}_B = \mathcal{C}_M \subseteq \mathcal{T}_M$.

The containment in the previous item is strict, when M is not idempotent. This is the case in example 2.

Lemma 7. *Let A be a NI matrix. Then the columns of A are pairwise different if and only if A is tropically regular.*

Proof. Of course, if A is tropically regular, then the columns of A are pairwise different. For the converse, if A is NI, then $|A|_{\text{trop}} = 0$ and this value is attained by the identity permutation. Suppose that A is tropically singular. Then there exists a permutation $\text{id} \neq \sigma$ such that $\sum_{j\in[n]} a_{j\sigma(j)} = 0$. Since each a_{ij} is non–positive, then $a_{j\sigma(j)} = 0$, for all $j \in [n]$.

Let us factor σ as a finite composition of disjoint cycles and let τ be one such cycle. By simplicity in writing, let us assume that $\tau = (1, 2, \ldots, s)$, with $2 \leq s \leq$

n. By hypothesis, $a_{12} = a_{23} = \cdots = a_{s-1s} = a_{s1} = 0$. Using $a_{ij} + a_{jk} \leq a_{ik} \leq 0$, we get $a_{ij} = 0$, for all $i, j \in [s]$. Thus, A has a $s \times s$ zero block. Now using $A = A^2$, we get $a_{ij} = a_{kj}$, for all $i, k \in [s]$, $j \in \{s+1, \ldots, n\}$ and $a_{ji} = a_{jk}$, for all $i, k \in [s]$, $j \in \{s+1, \ldots, n\}$. In particular, the first s columns of A are identical. In general, if $\tau = (i_1, \ldots, i_s)$, we obtain that the columns $i_1, \ldots, i_s$ of A are identical, with $s \geq 2$. ☐

Lemma 8. *If A is normal and tropically regular, then $\mathcal{C}_A = \mathcal{T}_{A^{n-1}}$.*

Proof. We know that A^{n-1} is NI.

Suppose that A^{n-1} has repeated columns. Then, there is at least one classical hyperplane Γ passing through all the columns of A^{n-1}. By (4), $\operatorname{col}(A^{n-1}, j) \in \overline{S_j}$ for all j, and we know that $\bigcap_{j \in [n]} \overline{S_j}$ is the origin. Then Γ goes through the origin. Since, for every $i, j \in [n]$, $i \neq j$, the direction of $\overline{S_i} \cap \overline{S_j}$ is a zero–one vector, then the direction of at least one hyperplane Γ is a zero–one vector. We conclude that the columns of A^{n-1} belong to the tropical hyperplane Π_0 given by expression (3) with $p = 0$. Thus, $\mathcal{T}_{A^{n-1}}$ is (an alcoved polytope) sitting inside Π_0. In p. 6 we recalled that $\mathcal{T}_A$ is connected and the difference $\mathcal{T}_A \setminus \mathcal{T}_{A^{n-1}}$ is a finite union of cells of dimension m, with $0 \leq m \leq n-2$. These cells are contained in classical hyperplanes $x_j = \operatorname{cnst}.$ or $x_j - x_k = \operatorname{cnst}.$, for some $j, k \in [n-1]$. Then, $\mathcal{T}_A$ is sitting inside Π_0 too and, in particular, A is tropically singular, a contradiction. Therefore, A^{n-1} has pairwise different columns. By lemma 7, A^{n-1} is tropically regular, so that $\mathcal{T}_{A^{n-1}}$ is a convex $n-1$–dimensional set. Moreover, $\mathcal{T}_{A^{n-1}}$ is maximal convex $n-1$–dimensional contained in $\mathcal{T}_A$ by [5, 9, 30]. Now, $\mathcal{T}_{A^{n-1}} = \mathcal{C}_{A^{n-1}} \subseteq \mathcal{C}_A \subseteq \mathcal{T}_A$, whence $\mathcal{C}_A = \mathcal{T}_{A^{n-1}}$, by maximality. ☐

In [28], $\mathcal{T}_{A^{n-1}}$ was called *soma* of $\mathcal{T}_A$, trying to express the fact that, in the generic case, $\mathcal{T}_{A^{n-1}}$ is the top dimensional part of $\mathcal{T}_A$.

5 Intersection of tropical simplexes defined by NI matrices

Theorem 9. *If A, B are NI, then A and B commute if and only if $\mathcal{T}_{AB} = \mathcal{T}_{BA} = \mathcal{T}_A \cap \mathcal{T}_B$. This is an alcoved polytope and a tropical simplex.*

Proof. By theorem 3 and item 1 in the previous list, we have $\mathcal{T}_{AB} = \mathcal{T}_{BA}$, from the start. Take $x \in \mathcal{T}_A \cap \mathcal{T}_B$. Then x is fixed by f_A and f_B, whence x is fixed by f_{AB}, proving that $x \in \mathcal{T}_{AB}$. Now take $x \in \mathcal{T}_{AB}$, so that $ABx = x$. Then, $Ax = A(ABx) = A^2Bx = ABx = x$, by idempotency of A. Thus, $x \in \mathcal{T}_A$. Similarly, $x \in \mathcal{T}_B$. ☐

Theorem 10. *If A, B are NI and M is tropically regular, then $\mathcal{T}_{M^{n-1}} = \mathcal{T}_A \cap \mathcal{T}_B$. This is an alcoved polytope and a tropical simplex.*

Proof. $\mathcal{T}_A = \mathcal{C}_A$ and $\mathcal{T}_B = \mathcal{C}_B$, whence $\mathcal{T}_A \cap \mathcal{T}_B = \mathcal{C}_A \cap \mathcal{C}_B = \mathcal{C}_M = \mathcal{T}_{M^{n-1}}$, by lemma 8 and item 2 in the previous list. ☐

Lemma 3.10 in [25] is somewhat related to our results in this section.

The same questions remain open for normal matrices A, B. Question 1: find conditions for A, B to commute under tropical multiplication. Question 2: is the intersection of the two tropical simplexes $\mathcal{T}_A, \mathcal{T}_B$ a tropical simplex?

References

[1] M. Akian, R. Bapat and S. Gaubert, *Max–plus algebra*, chapter 25 in *Handbook of linear algebra,* L. Hobgen (ed.) Chapman and Hall, 2007.

[2] X. Allamigeon, S. Gaubert and E. Goubault, *Computing the extreme points of tropical polyhedra*, ArXiv:0904.3436v3, 2011.

[3] F.L. Baccelli, G. Cohen, G.J. Olsder and J.P. Quadrat, *Syncronization and linearity,* John Wiley; Chichester; New York, 1992.

[4] P. Butkovič, *Simple image set of* $(\max, +)$ *linear mappings,* Discrete Appl. Math. **105**, (2000), 73–86.

[5] P. Butkovič, *Max–plus linear systems: theory and algorithms,* (2010), Springer.

[6] P. Butkovič, H. Schneider and S. Sergeev, *Generators, extremals and bases of max–cones*, Linear Algebra Appl. **421**, 394–406, (2007).

[7] R. Cuninghame–Green, *Minimax algebra*, LNEMS, **166**, Springer, 1970.

[8] R.A. Cuninghame–Green, *Minimax algebra and applications,* in *Adv. Imag. Electr. Phys.*, **90**, P. Hawkes, (ed.), Academic Press, 1–121, 1995.

[9] M. Develin, B. Sturmfels, *Tropical convexity*, *Doc. Math.* **9**, 1–27, (2004); Erratum in Doc. Math. **9** (electronic), 205–206, (2004).

[10] A. Gathmann, *Tropical algebraic geometry*, Jahresbericht der DMV, **108**, n.1, (2006), 3–32.

[11] S. Gaubert, R. Katz, *Max–plus convex geometry*, R.A. Schmidt (Ed.), RelMiCS/AKA 2006, LNCS 4136, (2006), 192–206.

[12] M. Gondran and M. Minoux, *Graphs, dioids and semirings. New models and algorithms*, Springer, 2008.

[13] J. Gunawardena (ed.), *Idempotency*, Publications of the Newton Institute, Cambridge U. Press, 1998.

[14] I. Itenberg, G. Mikhalkin and E. Shustin, *Tropical algebraic geometry*, Birkhäuser, 2007.

[15] Z. Izhakian, M. Johnson and M. Kambites, *Idempotent tropical matrices and finite metric spaces*, arXiv: 1203.2480, 2012.

[16] Z. Izhakian, M. Johnson and M. Kambites, *Pure dimension and projectivity of tropical politopes*, arXiv: 1106.4525, 2012.

[17] Z. Izhakian, M. Johnson and M. Kambites, *Tropical matrix groups*, arXiv: 1203.2449, 2012.

[18] A. Jiménez and M.J. de la Puente, *Characterizing the convexity of the n–dimensional tropical simplex and the six maximal convex classes in* $\mathbb{R}^3$, arXiv: 1205.4162, 2012.

[19] M. Joswig and K. Kulas, *Tropical and ordinary convexity combined*, Adv. Geom. **10**, (2010) 333-352.

[20] H.W. Kuhn, *The Hungarian method for the assignment problem*, Naval Res. Logist. **2**, (1955), 83–97.

[21] T. Lam and A. Postnikov *Alcoved polytopes I*, Discrete Comput. Geom., **38** n.3, 453-478 (2007).

[22] T. Lam and A. Postnikov *Alcoved polytopes II*, arXiv:1202.4015v1 (2012).

[23] G.L. Litvinov, V.P. Maslov, (eds.) *Idempotent mathematics and mathematical physics*, Proceedings Vienna 2003, American Mathematical Society, Contemp. Math. **377**, (2005).

[24] G.L. Litvinov, S.N. Sergeev, (eds.) *Tropical and idempotent mathematics*, Proceedings Moscow 2007, American Mathematical Society, Contemp. Math. **495**, (2009).

[25] G. Merlet, *Semigroup of matrices acting on the max–plus projective space*, Linear Algebra Appl. **432**, (2010), 1923–1935.

[26] G. Mikhalkin, *What is a tropical curve?*, Notices AMS, April 2007, 511–513.

[27] C.H. Papadimitriou and K. Steiglitz, *Combinatorial optimization: algorithms and complexity*, Prentice Hall, 1982 and corrected unabrideged republication by Dover, 1998.

[28] M. J. de la Puente, *Tropical linear maps on the plane*, Linear Algebra Appl. **435**, n. 7, (2011) 1681–1710.

[29] J. Richter–Gebert, B. Sturmfels, T. Theobald, *First steps in tropical geometry*, in [23], 289–317.

[30] S. Sergeev, H. Scheneider and P. Butkovič, *On visualization, subeigenvectors and Kleene stars in max algebra*, Linear Algebra Appl. **431**, 2395–2406, (2009).

[31] A. Werner and J. Yu, *Symmetric alcoved polytopes*, arXiv: 1201.4378v1 (2012).

[32] M. Yoeli, *A note on a generalization of boolean matrix theory*, Amer. Math. Monthly **68**, n.6, (1961) 552–557.

[33] U. Zimmermann, *Linear and combinatorial optimization in ordered algebraic structures*, Annals of discrete mathematics **10**, North–Holland, 1981.

J. Linde. Facultad de Matemáticas. Universidad Complutense. Madrid. Spain. `jorgelinde236@gmail.com`
M. J. de la Puente. Dpto. de Algebra. Facultad de Matemáticas. Universidad Complutense. Madrid. Spain. `mpuente@mat.ucm.es`


3
1
2
4
3
2
1
0
−1
−2
−3
Z
−4
−2
0
2
4
6
X
−3
−2
−1
0
1
2
3
Y